

READ SAFETY DIRECTIONS

Premise[®]

COCKROACH GEL

Active Constituent: 21.5 g/kg IMIDACLOPRID

GROUP 4A INSECTICIDE

For the control of cockroaches in the situations specified in the DIRECTIONS FOR USE

GENERAL INSTRUCTIONS

Apply using suitable applicator such as BAITGUN™. Gel should be applied out of reach of children, pets and other animals. Do not place gel on surfaces which are regularly washed.

Insecticide Resistance Warning

For insecticide resistance management PREMISE Cockroach Gel is a Group **4A** insecticide. Some naturally occurring insect biotypes resistant to PREMISE and other Group **4A** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if PREMISE or other Group **4A** insecticides are used repeatedly. The effectiveness of PREMISE on resistant individuals could be significantly reduced. Since occurrence or resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of PREMISE to control resistant insects. PREMISE may be subject to specific resistance management strategies. For further information contact your local supplier or your local Bayer Environmental Science representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Do **NOT** contaminate streams, rivers or waterways with the gel or used syringes.

STORAGE AND DISPOSAL

The syringe must not be sold separately. Do not destroy carton while product still remains.

Keep out of reach of children. Do not store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

LIMITATION OF LIABILITY

This product is designed only for the particular purposes indicated on this label. Failure to use this product strictly as directed may be illegal; prove dangerous; render the product ineffective and avoid any Warranty or Liability of Bayer Crop Science Pty Ltd.

Premise[®] is a trademark of Bayer

NRA Approval No. 52197/0899

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

Situations	Pest	Rate	Critical Comments
Barracks, Farm buildings, Food processing establishments, Factories, Homes, Hospitals, Meatworks (non- product areas), Offices, Restaurants, Schools, Shops, Storerooms, Transport vehicles & facilities.	<u>Small cockroaches</u> e.g. German Cockroach (<i>Blattella germanica</i>)	<u>Light infestations</u> 1 placement per m ² <u>Heavy infestations</u> 2 placements per m ² 1 placement = a 5 mm ball (about the size of a small pea or approx. 0.1 g) or a line about 2.5cm long	Apply gel in out of the way locations where cockroaches occur eg., under fridges, near water heaters, behind microwave ovens, garbage bins etc. Re-assessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist.
	<u>Large cockroaches</u> e.g. American Cockroach (<i>Periplaneta spp</i>)	<u>Light infestations</u> 2 placements per m ² <u>Heavy infestations</u> 3 placements per m ²	<i>Periplaneta</i> spp. require more bait because of the quantity which can be consumed by a single cockroach. Reassess baiting effectiveness as above.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Do not allow to contact food, livestock feed or their water. Do not place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.