

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



QuickBayt[®]

FLY BAIT

Active Constituents: 5.0 g/kg IMIDACLOPRID
1.0 g/kg (Z)-9-TRICOSENE

GROUP	4A	INSECTICIDE
-------	----	-------------

Ready-to-use fly bait granules for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTION FOR USE

QuickBayt contains BITREX, an acutely bitter agent which reduces the possibility of accidental ingestion by children, domestic animals or wildlife.

GENERAL INSTRUCTIONS

QuickBayt can be applied in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. For best results time applications to begin at the start of the season before fly populations have reached their peak.

Fly control should be considered part of an integrated strategy which incorporates good sanitation, appropriate management of breeding sites (eg. manure and other decaying organic matter) and exclusion measures for preventing access of flies into internal areas.

NOTE: This product is not attractive to all species of flies and control of non-target species may require other means of control.

Insecticide Resistance Warning

For insecticide resistance management, QuickBayt Fly Bait is a Group **4A** insecticide.

Some naturally occurring insect biotypes resistant to QuickBayt and other Group **4A** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if QuickBayt or other Group **4A** insecticides are used repeatedly. The effectiveness of QuickBayt on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of QuickBayt to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science or Bayer Animal Health representative.

PRECAUTIONS

Do NOT allow contact with raw or processed food, utensils or packaging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT place baits in locations accessible to domestic animals, livestock or birds. Do NOT contaminate streams, rivers or waterways with the chemical or used containers. Do not scatter bait on manure heaps or dispose of treated sacks/cardboard/fabric with manure. Clean up spillage promptly. Very highly toxic to bees.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Disposal - 350 g Pack:

Shake container empty when completing use and dispose of empty container by wrapping with paper and putting in the garbage.

Disposal - 2 Kg pack:

Triple or preferably pressure rinse containers before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit. Empty containers and product should not be burnt.

Treated sacks, cardboard or fabric may be re-used as necessary. After use they should be disposed of by wrapping in paper, placing in a plastic bag and putting in the garbage.

SAFETY DIRECTIONS

Will irritate eyes. Avoid contact with eyes. If product in eyes wash it out immediately with water. Wash hands after use. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and rubber gloves. After each days use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 03 4747000).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No. 57892/0704

QuickBayt® is a Registered Trademark of Bayer.

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE**Restraint**

DO NOT place bait in areas accessible to food producing animals or pets.

PEST	Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)
RATE	2.0 g/m ²
SITUATION	CRITICAL COMMENTS
<p>Outside: Abattoirs, animal housing, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around: Industrial sites, feedlot manure heaps, garbage dumps</p>	<p>Scatter application: The bait should be scattered (do not put in piles) onto dry level surfaces (eg. pathways, window ledges or in shallow dishes). Avoid extremely dusty areas.</p> <p>Bait Station application Set up bait stations by sprinkling into shallow tins, lids or similar containers and place where flies congregate. Position bait stations out of reach of animals or livestock and away from feed troughs, storage or mixing areas.</p> <p>Wet Bait: In many situations more flies can be attracted to moist bait. This can be achieved by sprinkling bait on moist hessian sacks (placed out of reach of all livestock) or by lightly moistening bait with a light spray of water.</p> <p>Paint-on application: To prepare a paste for painting, mix 50 g of QuickBayt with 37.5 mL of water. 15 minutes after mixing the paste can be applied with a brush to spots where flies rest (eg. Walls, joists, window ledges, stable fixtures). Additionally it can be painted onto hessian bags or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>Re-baiting: Re-baiting may be necessary in 3-5 days depending on exposure, weather and density of fly populations.</p>
<p>Inside: Commercial and industrial facilities and animal houses ONLY in areas inaccessible to livestock and domestic animals</p> <p>Under: Animal and poultry cages ONLY in areas inaccessible to livestock</p>	<p>Apply as a scatter bait, in bait stations or as a paint-on application as above.</p> <p>Place out of reach of all livestock and away from feed.</p> <p>Re-bait dry bait stations as above.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.