

500 g pouch front panel

---

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS**



**Racumin<sup>®</sup>**  
**Rat And Mouse Paste**

**ACTIVE CONSTITUENT: 0.37 g/kg COUMATETRALYL**

Ready-to-use baits in a palatable sachet for the control of rats and mice

**Contains 50 bait sachets which it is illegal to sell separately.**

*(rat pictogram)*

500 g NET (50 x 10g bait sachets)

## DIRECTIONS FOR USE

**Restraints - Do NOT place baits in open unless in bait stations.**

**Do NOT place baits in any position accessible to children, livestock, or domestic pets.**

**Do NOT remove bait from the paper sachet.**

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, industrial, wharf areas, around farm buildings, dairies, meatworks (non-product areas)	Mice and Rats	3-6 sachets per bait site	Place bait sachets in areas of known rodent activity. For rats place a minimum of 3 sachets (30g) per placement. Do not exceed 3 m between baits for mice, or 9 m for rats. Follow the <b>Baiting Strategy</b> detailed below.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS:** Do not apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, food storage areas, or onto packaging materials or in areas where such packaging is stored. Do not feed the bait to wild or domestic birds.

### Baiting Strategies

Eliminate, as far as practicable, all alternative food sources. Place the bait sachets in areas of known rodent activity and under cover (eg. Behind appliances and furniture, in ceiling voids etc). Inspect baits daily. If eaten quickly increase the number of bait sachets. Continue observation and replenishment until no more bait is taken. Baiting for at least 2 weeks is necessary to reduce mouse or rat numbers. Caution is required in use of this product in and near piggeries. Do NOT allow to contaminate pig food.

**PRECAUTION:** Should not be used in areas accessible to children.

### PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Place baits in locations inaccessible to other animals and birds. This product is hazardous to pigs, cats and dogs. Do not contaminate ponds, waterways or drains with the product, bait, refuse from the bait stations, or used containers.

500 g pouch back panel

---

**STORAGE AND DISPOSAL:**

Store in the tightly closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight. Destroy unopened baits. Any spillage of bait should be removed immediately. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

**SAFETY DIRECTIONS:**

Should not be used in areas accessible to children. Poisonous if swallowed. If on skin, and after each baiting, wash thoroughly with soap and water.

**FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126. Vitamin K1 (Phytomenadione) is antidotal.

**MATERIAL SAFETY DATA SHEET:**

Additional information is listed in the Material Safety Data Sheet.


**Exclusion of Liability**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Bayer Environmental Science  
A Business Group of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn, Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)

APVMA Approval No. 51508/500 g/0704

[B]  
[DOM]



Racumin® is a trademark of Bayer  
Bitrex™ is a trademark of Macfarlan Smith Limited

5 kg bucket front panel

---

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS**



**Racumin<sup>®</sup>**  
**Rat And Mouse Paste**

**ACTIVE CONSTITUENT: 0.37 g/kg COUMATETRALYL**

Ready-to-use baits in a palatable sachet for the control of rats and mice

**Contains 50 bait sachets which it is illegal to sell separately**

*(rat pictogram)*

**5 kg NET (50 x 100 g bait sachets)**

## DIRECTIONS FOR USE

**Restraints - Do NOT place baits in open unless in bait stations.**

**Do NOT place baits in any position accessible to children, livestock, or domestic pets.**

**Do NOT remove bait from the paper sachet.**

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, industrial, wharf areas, around farm buildings, dairies, meatworks (non-product areas)	Mice and Rats	1 sachet per bait site	Place bait sachets in areas of known rodent activity. Do not exceed 3 m between baits for mice, or 9 m for rats. Follow the <b>Baiting Strategy</b> detailed below.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS:** Do not apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, food storage areas, or onto packaging materials or in areas where such packaging is stored. Do not feed the bait to wild or domestic birds.

### **Baiting Strategies**

Eliminate, as far as practicable, all alternative food sources. Place the bait sachets in areas of known rodent activity and under cover (eg. Behind appliances and furniture, in ceiling voids etc). Inspect baits daily. If eaten quickly increase the number of bait sachets. Continue observation and replenishment until no more bait is taken. Baiting for at least 2 weeks is necessary to reduce mouse or rat numbers. Caution is required in use of this product in and near piggeries. Do NOT allow to contaminate pig food.

**PRECAUTION:** Should not be used in areas accessible to children.

### **PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:**

Place baits in locations inaccessible to other animals and birds. This product is hazardous to pigs, cats and dogs. Do not contaminate ponds, waterways or drains with the product, bait, refuse from the bait stations, or used containers.

5 kg bucket back panel

---

**STORAGE AND DISPOSAL:**

Store in the tightly closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight. Destroy untaken baits. Any spillage of bait should be removed immediately. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

**SAFETY DIRECTIONS:**

Should not be used in areas accessible to children. Poisonous if swallowed. If on skin, and after each baiting, wash thoroughly with soap and water.

**FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126. Vitamin K1 (Phytomenadione) is antidotal.

**MATERIAL SAFETY DATA SHEET:**

Additional information is listed in the Material Safety Data Sheet.

**Exclusion of Liability**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Bayer Environmental Science  
A Business Group of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn, Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)

APVMA Approval No. 51508/5kg/0704

[B]  
[DOM]

APN Barcode

Racumin® is a Registered Trademark of Bayer  
Bitrex™ is a trademark of Macfarlan Smith Limited

'Tea-bags' – both sizes

---

# **Racumin<sup>®</sup> Paste**

**Coumatetralyl 0.375 g/kg**

