

**DIRECTIONS FOR USE**

**Restraints:**  
DO NOT apply bait to crops  
DO NOT place bait in the open  
DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments

**Baiting Strategy**

- Eliminate as far as practical, all alternative food sources
- Bait infested area, place baits under cover
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used
- Ensure bait blocks are removed when rodent activity has ceased

**Clean-Out Treatments (Pulse Baiting)**

- Clean-Out treatments are used to remove existing infestations of rats and mice
- Always record the number and locations of bait blocks for the purpose of bait replacement and later removal
- For rats, place 1 to 3 blocks at intervals of 5 to 9 m in infested areas
- For mice, place single blocks at 2 to 3 m intervals in infested areas
- Nails or wire may be used to fix blocks to prevent them being removed by rodents. Alternatively, baits may be secured in bait stations
- Bait stations should be secured to non-target animals. Use tamper-resistant bait stations for monitoring and maintenance
- Use the greater number of blocks and shorter distance between bait placements when infestations are heavy
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rodent numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- After the infestation has been controlled, a regular interval schedule and remove all bait and bait stations unless a maintenance treatment is to be undertaken

**Monitoring and/or Maintenance Treatments**


Monitoring and/or maintenance treatments are required at sensitive sites where there is a history of rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatments may be a condition for audit

approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.

- Always record the number and locations of bait stations for the purpose of bait replacement and later removal
- For rats and mice place one block at intervals of between 15 and 30 m in areas prone to rodent infestation
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance
- Bait stations should be secured to non-target animals
- Use the greater number of blocks and shorter distance between bait placements when infestations are heavy
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rodent numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- After the infestation has been controlled, a regular interval schedule and remove all bait and bait stations unless a maintenance treatment is to be undertaken

Application	Situation	Pests	Rate	Critical Comments
	Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	Rats ( <i>Rattus</i> 1 to 3 blocks/ <i>rattus</i> , <i>R. norvegicus</i> ) Mice ( <i>Mus domesticus</i> )	Station	Set out bait in places frequented by mice and rats in and around buildings, fencelines (within 2 m) or enclosed spaces, eg drains. Clean-Out Treatment: For rats: Place 1 to 3 blocks at intervals of 5 to 9 m in infested areas. For mice: Place single blocks at 2 to 3 m intervals in infested areas. See also Monitoring and/or Maintenance Treatments.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**



**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**MAX BLOCKS**

**Talona**  
RODENTICIDE

CONTAINS BITECIX

**ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUm**

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, public services, agricultural and domestic buildings. Controls rodents resistant to other anticoagulants, including warfarin, coumatetralyl and bromadiolone

**ONE FEED KILLS RATS AND MICE\***

1 kg NET

Syngenta Crop Protection Pty Limited  
Level 1, 2-4 Lygonpark Road, Macquarie Park NSW 2113  
In a transport emergency dial 000, Police or Fire Brigade  
For specialist advice in an emergency only.  
APVMA Approval No. 52675/1/0109  
TINW81A/02

**syngenta**

U036 454x115 FLAT die

**GENERAL INSTRUCTIONS**

Ensure baits are inaccessible to children, pets, domestic animals and wildlife. Place bait along walls or rafters and in dark, sheltered spots where there are signs of rodent activity. Nail in place if relevant. Wax blocks are particularly suitable for use in damp situations and sewers, and crop stores where the risk of contamination should be minimised. Record the number and location of bait placements for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Look for dead rats or mice, burn or bury them. DO NOT place in ethane bins or rubbish tips. Ensure bait blocks are secured and covered with rodent proof caps. Ensure bait blocks are #Note: One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimize this.

**PRECAUTION**

Avoid contact with food, food utensils, or places where food is prepared or stored.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. DO NOT place baits in locations which are accessible to pets, domestic animals, livestock or birds. DO NOT contaminate dams, waterways and drains with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities. The product is restricted to use in and around buildings, fencelines (within 2 m) or enclosed spaces, eg drains.

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If no landfill is available, bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

**SAFETY DIRECTIONS**

Poisonous if swallowed. DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. Vitamin K1 (Phytomenadione) is antidotal.

**NOTE TO PHYSICIANS AND VETERINARIANS**

Brodifacoum, the active constituent of TALON Wax Blocks, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. It is not an anti-emetic and the emetic properties of ipecacuanha may be ineffective against the emetogenic response of veterinarians. If administered intravenously the injection must be given slowly.

**MATERIAL SAFETY DATA SHEET**

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108, or visit our website at www.syngenta.com.au

**MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Syngenta has no control over storage, handling and use of this product. It is sold as a baiting product and is intended to be used correctly and in accordance with directions. No express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

\* Product names marked @, the SYNGENTA Logo and the CP FRAME are trademarks of a Syngenta Group Company  
#Bitecix is a trademark of Mactarlan Smith Limited

D.O.M.	Batch No.

91322642001407

595.23

APPROVED LABEL

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Food safety guidelines. However, all use of this product must comply with the conditions of this label.

**Directions for Use**

Retraite  
DO NOT apply bait to crops  
DO NOT place bait in the open  
DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments

**Baiting Strategy**

- Eliminate as far as practical, all alternative food sources
- Bait infested area, place baits under cover.
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used
- Ensure bait blocks are removed when rodent activity has ceased

**Clean-Out Treatments (Pulse Baiting)**

- Clean-Out treatments are used to remove existing infestations of rats and mice
- Always record the number and locations of bait blocks for the purpose of bait replacement and later removal
- For rats, place 1 to 3 blocks at intervals of 5 to 9 m in infested areas
- Nails or wire may be used to fix blocks to prevent them being removed by rodents. Alternatively, baits may be secured in bait stations.
- Place baits so as to prevent access to non-target animals. Use tamper-resistant bait stations at locations where access of non-target animals cannot otherwise be prevented and fix the stations securely to the substrate, where necessary.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any placements when infestations are heavy
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Monitor and maintenance treatments are required at sensitive sites and remove all bait and bait stations unless a maintenance treatment is to be undertaken

**Monitoring and/or Maintenance Treatments**

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatments may be a condition for audit approval within

Situation	Pests	Bait	Critical Comments
Industrial, Commercial, Public, Services, Agricultural and Domestic Buildings	Rats ( <i>Rattus norvegicus</i> ), Mice ( <i>Mus domesticus</i> )	1 to 3 bait stations	Set out bait in places frequented by rodents, eg buildings, enclosed spaces, eg drains. Clean-Out Treatment: For rats: Place 1 to 3 blocks at intervals of 5 to 9 m in infested areas. For mice: Place single blocks at 2 to 3 m intervals in infested areas. See also Monitoring and/or Maintenance Treatments.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**MAX BLOCKS**

**Talor®**  
RODENTICIDE

**BiteK**  
CONTAINS VITAMIN K1  
ANTICOAGULANT

**ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM**

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, public services, agricultural and domestic buildings. Controls rodents resistant to other anticoagulants, including warfarin, coumatetralyl and bromadiolone

**ONE FEED KILLS RATS AND MICE#**

**2.4 kg NET**

Syngenta Crop Protection Pty Limited  
Level 1, 2-4 Lygonpark Road, Macquarie Park NSW 2113  
In a transport emergency dial 000, Police or Fire Brigade  
For specialist advice in an emergency only,  
call 1800 053 111 (24 hours)  
Australia 4 8194 4400  
T8W82 444U02

**syngenta®**

**GENERAL INSTRUCTIONS**

Ensure baits are inaccessible to children, pets, domestic animals and wildlife. Place bait along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Bait in place if relevant. Wax or grease bait stations to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.

Record the number and location of bait placements for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Look for dead rats or mice, turn out any items, for example, food, bedding, etc. Ensure bait blocks are removed when rodent activity has ceased.

**#Note:** One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimize this.

**PRECAUTION**

Avoid contact with food, food utensils, or places where food is prepared or stored.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Depending on the amount eaten, this bait may kill or injure the bait is baits in locations which are accessible to pets, domestic animals, livestock or birds. DO NOT contaminate dams, waterways and drains with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from the relevant authority. Do not use this product in or around buildings, structures, fences (within 2 m) or enclosed spaces, eg drains.

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If no landfill is available, bury the containers or unused bait in a secure container. Do not use this product in or around buildings, structures, fences (within 2 m) or enclosed spaces, eg drains. Empty containers and product should not be burnt. DO NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

**SAFETY DIRECTIONS**

Poisonous if swallowed, DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. Vitamin K1 (Phylloquinone) is antidotal.

**NOTE TO PHYSGIANS AND VETERINARIANS**

Brodifacoum, the active constituent of TALON Max Blocks, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phylloquinone) should be administered urgently by approved appropriate personnel. The antidote should be administered intravenously. The injection must be given slowly.

**MATERIAL SAFETY DATA SHEET**

Material Safety Data Sheet is required refer to the Material Safety Data Sheet For a copy phone 1800 057 108, or visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

**MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Syngenta warrants that the material is not forged, handled or used contrary to the directions, and in accordance with directions, no express or implied warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

© Product names marked ®, the SYNGENTA Logo and the CP FRAME are trademarks of a Syngenta Group Company  
BiteK is a trademark of Macfarlan Smith Limited

Batch Number

Date of Manufacture

913 2 26 420 01 3911

**Recyclable container**